

## TSA - Reverse Arthroplasty – Postoperative Instructions

### DIET

- Begin with clear liquids and light foods (jellos, soups, etc.).
- Progress to your normal diet if you are not nauseated.

### WOUND CARE

- Maintain your operative dressing, loosen bandage if swelling of the hand occurs.
- It is normal for the shoulder to bleed and swell following surgery. If blood soaks through the bandage, do not become alarmed, reinforce with additional dressing.
- To avoid infection, keep surgical incisions clean and dry for the first **7 days** following surgery – you may shower by placing a large plastic bag over your sling beginning the day after surgery. NO immersion of the operative shoulder (ie: bath or pool).
- Wait until your first post operative appointment to have Dr. Chahla's team remove the surgical dressing.
- Please do not place any ointments lotions or creams directly over the incisions.
- Once the sutures/staples are removed **at least 14 days post operatively** you can begin to get the incision wet in the shower (water and soap lightly run over the incision and pat dry).
- NO immersion in a bath until given approval by our office.

### MEDICATIONS

- Local anesthetics are injected into the wound and shoulder joint at the time of surgery. This will wear off within 8-12 hours and it is not uncommon for patients to encounter more pain on the first or second day after surgery when swelling peaks.
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per directions on the bottle.
- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease the side effects take the medication with food. If constipation occurs, consider taking an over the counter laxative.
- If you are having problems with nausea and vomiting, contact the office to possibly have your medications changed.
- Do not drive a car or operate machinery while taking the narcotic medication or while in sling.
- Please avoid alcohol use while taking narcotic pain medication.
- If you are having pain that is not being controlled by the pain medication prescribed, you may take an over the counter anti-inflammatory medication such as ibuprofen or naproxen in between doses of pain medication. This will help to decrease pain and decrease the amount of narcotic medication required. Please take as directed on the bottle.
- For 2 weeks following surgery take one aspirin daily to lower the risk of developing a blood clot after surgery. Please contact the office should severe calf pain occur or significant swelling of the calf or ankle occur.
- 3-5 day course of antibiotics will be prescribed post operatively to decrease the chance of infection.
- Please call if planning on having dental procedure done in the future as you will also need antibiotics prior to future dental procedures.

### ACTIVITY

- You are to wear the sling placed at surgery for a total of 6 weeks as described by Dr. Chahla. This includes sleeping and throughout the day.
- If there are 24 hours a day, you should be in the sling 23.5 hours of the day. Removal for hygiene, dressing, and home exercise only.
- When sleeping or resting, inclined positions (ie: reclining chair) and a pillow under the forearm for support may provide better comfort **STILL IN SLING**.

- Do not engage in activities which increase pain/swelling. Unless otherwise instructed the arm should remain in the sling at all times.
- Avoid long periods of sitting or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician, it is illegal to drive in a sling.
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable.

#### ICE THERAPY

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Use icing machine continuously or ice packs (if machine not prescribed) for 30-45 minutes every 2 hours daily until your first post-operative visit.
- Care should be taken with icing to avoid frostbite to the skin.
- You do not need to wake up in the middle of the night to change over the ice machine or icepacks unless you are uncomfortable.

#### EXERCISE

- Begin exercises (active elbow flexion/extension without resistance) 24 hours after surgery unless otherwise instructed.
- While maintaining your elbow by the side, begin elbow, hand, and wrist exercises immediately.
- Formal physical therapy (PT) typically begins 2-4 weeks following surgery.

#### EMERGENCIES\*\*

Contact Dr. Chahla's team at [chahlaadmin@rushortho.com](mailto:chahlaadmin@rushortho.com) if any of the following are present:

- Painful swelling or numbness (note that some swelling and numbness is normal).
- Unrelenting pain.
- Fever (over 101° – it is normal to have a low grade fever for the first day or two following surgery) or chills.
- Redness around incisions.
- Color change in foot or ankle.
- Continuous drainage or bleeding from incision (a small amount of drainage is expected).
- Difficulty breathing.
- Excessive nausea/vomiting.
- Calf pain.
- If you have an emergency after office hours or on the weekend, contact the office at 312-432-2390 and you will be connected to our pager service. This will connect you with the Physician on call. You can also call Rush University Medical Center at 312-942-5000 and ask for the operator to page the orthopedic resident on call.
- If you have an emergency that requires immediate attention proceed to the nearest emergency room.

#### FOLLOW-UP CARE/QUESTIONS

- If you do not already have a post-operative appointment scheduled, please contact the our scheduler at 708-236-2701 to schedule.
- Typically the first post-operative appointment following surgery is 10-14 days following surgery.
- The first post operative appointment will be with one of the Physician Assistants. They will assess the wound, go over post operative protocol, and answer any questions you may have regarding the procedure.
- If you have any further questions please contact Dr Chahla's team at [chahlaadmin@rushortho.com](mailto:chahlaadmin@rushortho.com)