



Anterior Cruciate Ligament (ACL) Repair

Post-Operative Instructions

Post Operative Prescribed Medications:

- Aspirin: Take as prescribed for 30 days after surgery
- Norco (5-325mg): Take 1-2 tablets every 4-6 hours as needed for pain.
- Zofran (4mg): Dissolve 1 tablet on tongue every 8 hours as needed for nausea.

Physical Therapy:

- Begin on post op day 1 or 2. <u>Bring protocol</u> and order provided on surgery day to initial PT visit.
- Attend PT 2-3 times per week for the first 2-3 months after surgery
- Expect to be in PT for 6-8 months following surgery.

First Post-op Clinic Appointment:

 Your first post-op clinic visit will be 1-2 days after surgery.

Weight bearing after surgery:

 Your weightbearing status following surgery will rely on whether or not a meniscus repair is performed in conjunction with you ACL reconstruction. Your weightbearing status will be clearly outlined to you prior to leaving the surgery center.

Smoking, marijuana, and surgery:

- Dr. Chahla has a <u>NO NICOTINE</u> policy for surgical patients.
- For your safety and to promote successful outcomes, we recommend sustaining from marijuana use for 4 weeks post op. This includes oral, topical, and inhalation.

Dressings and Showering:

- Remove gauze and padded dressings on post op day 3.
- Please leave the steri-strips on until initial post-op appointment with one of Dr. Chahla's PAs.
- Do not shower until post op day 3. Beginning post op day 3, you can shower while keeping the incisions covered and dry. Do not submerge the incisions or use topical ointment for at least 4 weeks after surgery or until insicion is fully healed.

Return to Driving:

 You can return to driving once you are off crutches, no longer taking narcotics, and feel that you can come to an emergency stop.
This is typically around 1-8 weeks post op depending on if a meniscus repair is completed (less time if no repair)

Return to Work/School:

 You can expect to return to desk work and/or school about 4-7 days after surgery.



^{*}Please Note: medications may vary based off your current mediction list. These variations will be discussed during your pre-operative phone consultation with a PA.*