



Patella Open Reduction Internal Fixation (ORIF)

Post-Operative Instructions

Post Operative Prescribed Medications:

- Lovenox: Take as prescribed for 14 days after surgery.
- Aspirin: Take as prescribed for 14 days after completion of Lovenox
- Norco (5-325mg): Take 1-2 tablets every 4-6 hours as needed for pain.
- Keflex: Take as prescribed for 7 days after surgery.
- Zofran (4mg): Dissolve 1 tablet on tongue every 8 hours as needed for nausea.

Please Note: medications may vary based off your current medication list. These variations will be discussed during your pre-operative phone consultation with a PA.

Physical Therapy:

- Begin on post op day 1 or 2. Bring protocol and order provided on surgery day to initial PT visit.
- Attend PT 2-3 times per week for the first 2-3 months after surgery.
- Expect to be in PT for 4-6 months following surgery.

First Post-op Clinic Appointment:

- Your first post-op clinic visit will be 1-2 days after surgery unless instructed otherwise by a member of Dr. Chahla's Team.

Weight bearing after surgery:

- You will be weightbearing as tolerated by pain following surgery.

Smoking, marijuana, and surgery:

- Dr. Chahla has a NO NICOTINE policy.
- For your safety and to promote successful outcomes, we recommend sustaining from marijuana use for 4 weeks post op. This includes oral, topical, and inhalation.

Dressings and Showering:

- Remove gauze and padded dressings on post op day 3.
- Please leave the steri-strips on until initial post-op appointment with one of Dr. Chahla's PAs.
- Do not shower until post op day 3. Beginning post op day 3, you can shower while keeping the incisions covered and dry. Do not submerge the incisions or use topical ointment for at least 4 weeks after surgery or until incision is fully healed.

Return to Driving:

- You can return to driving once you are off crutches and brace, no longer taking narcotics, and feel that you can come to an emergency stop. This is typically around 6 weeks post op.

Return to Work/School:

- You can expect to return to desk work and/or school about 4-7 days after surgery.