

# Patellofemoral Arthroplasty Post-Operative Rehabilitation Protocol

Jorge Chahla, MD, PhD

#### 0-3 weeks

- WBAT with knee locked in extension
- Gentle PROM
- ROM goal by week 2: 0° of extension and 110° of flexion
- Progression off use of assistive device to unassisted ambulation beginning POD 0, guided by PT
- Heel slides
- Light strengthening of abductor group
- SLR supine with brace locked at 0°
- Quad Sets with brace locked at 0°
- Ankle Pumps

#### >3 weeks

- Unlock brace for ambulating, wean off brace once good quad control
- WBAT without assistive device
- Progress to full ROM
- Advance quad and abductor strengthening

Weight **Bearing Status** 

**WBAT** 

**Brace Settings** 

ambulating or

**ROM** Restrictions

> Full PROM POD 1



Exercise	Week													
	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Initial Exercises:	l l	<u> </u>			ı				ı					
ROM Goals	0-	0-110° Progress									I			
Flexion/extension, heel slides														
Flexion/extension- seated														
Patella/Tendon mobilization														
Quad Series														
Hamstring Sets														
Sit and Reach for Hamstrings														
Ankle Pumps														
Crutch weaning	W	BAT												
SLR (w/ brace until quad control restored)														
Toe and Heel Raises														
Balance Series														
Weight-bearing Strength Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Double Knee Bends		•												
Double Leg Bridges														
Reverse Lunge, static holds														
Beginning cord exercises														
Balance Squats														
Dead Lift (2 →1 leg)														
Leg Press (ISO →reps, 2 → 1 leg) (80-0° arc)														
Sports Test Exercises														
Cardiovascular Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Bike with both legs-no resistance									•	•				
Bike with both legs- resistance														
Aqua jogging														
Treadmill-walking 7% incline														
Swimming with fins-light flutter kick														
Elliptical Trainer														
Rowing														
Stair stepper														
Agility Exercises:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Running Progression														
Initial-Single Plane														
Advance-Multi Directional														
Functional Sports Test														
High Level Activities:	1	2	3	4	5	6	7	8	9	10	12	16	20	24
Golf Progression														
Outdoor biking, hiking, running														
Return to Full Sport														



#### **Quad Series:**

- •Quad sets (5 sec hold) also quick contractions (NMES to supplement)
- Prone quad sets
- •SLR with no extension leg
- Progression of Quad Strengthening
- •LAQ in safe range -no added resistance to begin
- •Once weight bearing tandem stance TKE with 5 sec hold
- •Isometric wall sit
- SLR with resistance
- •Step up progression
- Squat progression
- •Leg press progression
- Lunge progression

#### **Hamstring Series:**

- Hamstring stretching (supine/prone)
- •Hamstring set (heel dig)- contract hamstring with knee flexed pressing heel into the floor (5 sec hold)
- Hamstring curls
- Progress to resisted hamstring curl

#### **Sport Test Exercises:**

- •Goals: impact training and introduction of directional drills
- Double leg squat jumps
- Box jumps
- Speed skaters
- Double leg jumps for height and distance progress to single leg as tolerated
- •Blaze pod change of direction drills

#### Further Clarifications and Considerations

- •Double Knee Bends: feet shoulder width apart, bend knees to 30°, keeping knees behind the toes.
- •Cord exercises (resistance band/cord): resisted double knee bends, resisted hamstring curls
- •May begin to implement BFR as early as 1 week into post-op period



### Return to Running Criteria

- Trace effusion, flexion within 5 degrees of contralateral side
- LSI on anterior reach Y balance test ≥ 90%
- LSI on quadriceps torque output on isometric measurement ≥ 75%
- •12" SL squat tolerance
- Able to walk > 1 mile with no limping or pain
- Able to "hop" with UE assist on operative leg 5 times without pain or compensation
- •SL balance with eyes closed 30 seconds without loss of balance



# Return to Agility/Change of Direction Criteria

- •Return to running criteria listed above met
- No effusion
- Full ROM
- •Quad LSI on isokinetic ≥85%
- Hamstring LSI on isokinetic ≥85%
- •LSI on anterior reach Y-balance ≥ 95%
- SL hopping pain-free and effusion free



## Return to Sport Criteria

- •In order to test must demonstrate full AROM, no pain, no/trace effusion, 5/5 strength, negative lachman and pivot shift, LSI ≥95% hamstring curl and leg press
- •Quadricep strength ≥95% of contralateral side (< 2cm of atrophy)
- •Able to perform single leg squat with correct form
- •LSI SL hop ≥95%
- •Y-Balance ≥95% (mean of 3 trials in anterior, posterolateral, posteromedial /100)
- Vertical jump test, single leg hop distance, and timed single leg hop over 20 -feet ≥90% of contralateral

