



Posterolateral Corner Reconstruction

Post-Operative Therapy Protocol

0-2 weeks

- NWB for 6 weeks
- ROM: 0-90° with emphasis on full extension
- Patella mobilization
- SLR supine with brace locked at 0°
- Quad Sets with brace locked at 0°
- Ankle Pumps
- Short crank (90mm) ergometry

2-6 weeks

- Flexion extension wall slides

6-12 weeks

- Begin spinning on a stationary bike
- Begin to wean off crutches
- Sit and reach for hamstrings
- Progressive Squat program
- Initiate Step-Down program
- Balance Series
- Double leg bridges
- Begin cord exercise

12-20 weeks

- Sports test exercises
- No gravity running

>24 weeks



- Running progression
- Agility exercises
- Multidirectional motions
- Lower Body Assessment when initiating return to sport progression. Schedule via QR code email or spc@rushortho.com

Weight Bearing Status

NWB x 6 weeks

Brace Settings

Locked straight x 6 weeks when ambulating or sleeping. Otherwise open brace once good quad control

ROM Restrictions

Full PROM beginning POD 1



Exercise	Week																
	1	2	3	4	5	6	7	8	9	10	12	16	20	24			
Initial Exercises:																	
ROM Goals	0-90°		0-125°			Progress to full											
Flexion/extension, heel slides																	
Flexion/extension- seated/ calf-assisted or prone																	
Patella/Tendon mobilization																	
Quad Series	0° ISOs, SLR				60° ISOs			70-30° arc-resisted				Full arc resisted					
Hamstring Sets																	
Sit and Reach for Hamstrings																	
Ankle Pumps																	
Crutch weaning	NWB																
Heel prop knee extension stretch																	
SLR (w/ brace until quad control restored)																	
Toe and heel raises																	
Weight-bearing Strength Exercises:																	
Double Knee Bends	1	2	3	4	5	6	7	8	9	10	12	16	20	24			
Double Leg Bridges	NWB																
Step up/Lunge Progression																	
Beginning cord exercises																	
Proprioception/Balance Training																	
Dead Lift (2 → 1)																	
Squat/Leg Press (ISO →reps, 2→1 leg)																	
Sports Test Exercises																	
Cardiovascular Exercises:																	
Bike with both legs-no resistance	1	2	3	4	5	6	7	8	9	10	12	16	20	24			
Bike with both legs- resistance	NWB																
Aqua jogging																	
Treadmill-walking 7% incline																	
Swimming with fins-light flutter kick																	
Elliptical Trainer																	
Rowing																	
Stair stepper																	
*Cardio Exercises							Must tolerate daily walking with no limp & no significant increase in pain/swelling prior to initiating WB cardio (walk, elliptical, stepper)										
Agility Exercises:																	
Running Progression*	1	2	3	4	5	6	7	8	9	10	12	16	20	24			
Initial-Single Plane	NWB																
Advance-Multi Directional																	
Functional Sports Test																	
High Level Activities:																	
Golf Progression	1	2	3	4	5	6	7	8	9	10	12	16	20	24			
Outdoor biking, hiking, running	NWB																
Return to Full Sport at 9 months post-op																	



Quad Series:

- Quad sets (5 sec hold) also quick contractions (NMES to supplement)
- Prone quad sets
- SLR with no extension leg
- Progression of Quad Strengthening (WB exercises at 6 weeks)
 - LAQ in safe range -no added resistance to begin
 - Once weight bearing tandem stance TKE with 5 sec hold
 - Isometric wall sit
 - SLR with resistance
 - Step up progression
 - Squat progression
 - Leg press progression
 - Lunge progression

Hamstring Series:

- Hamstring stretching (supine/prone)
- Hamstring set (heel dig)- contract hamstring with knee flexed pressing heel into the floor (5 sec hold)
- Hamstring curls
- Progress to resisted hamstring curl

Sport Test Exercises:

- Goals: impact training and introduction of directional drills
- Double leg squat jumps
- Box jumps
- Speed skaters
- Double leg jumps for height and distance progress to single leg as tolerated
- Blaze pod change of direction drills


Further Clarifications and Considerations

- May begin to implement BFR as early as 1 week into post-op period
- Functional Sports Test: Baseline completed at 7 months post-op with subsequent test at 9 months




Return to Running Criteria

- Trace effusion, flexion within 5 degrees of contralateral side
- LSI on anterior reach Y balance test $\geq 90\%$
- LSI on quadriceps torque output on isometric measurement $\geq 75\%$
- 12" SL squat tolerance
- Able to walk > 1 mile with no limping or pain
- Able to "hop" with UE assist on operative leg 5 times without pain or compensation
- SL balance with eyes closed 30 seconds without loss of balance



Return to Agility/Change of Direction Criteria

- Return to running criteria listed above met
- No effusion
- Full ROM
- Quad LSI on isokinetic $\geq 85\%$
- Hamstring LSI on isokinetic $\geq 85\%$
- LSI on anterior reach Y-balance $\geq 95\%$
- SL hopping pain-free and effusion free



Return to Sport Criteria

- In order to test must demonstrate full AROM, no pain, no/trace effusion, 5/5 strength, negative lachman and pivot shift, LSI $\geq 95\%$ hamstring curl and leg press
- Quadricep strength $\geq 95\%$ of contralateral side (< 2cm of atrophy)
- Able to perform single leg squat with correct form
- LSI SL hop $\geq 95\%$
- Y-Balance $\geq 95\%$ (mean of 3 trials in anterior, posterolateral, posteromedial/100)
- Vertical jump test, single leg hop distance, and timed single leg hop over 20 -feet $\geq 90\%$ of contralateral side

